

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 : C07J 71/00	A1	(11) International Publication Number: WO 91/04984 (43) International Publication Date: 18 April 1991 (18.04.91)
(21) International Application Number: PCT/SE90/00619		(74) Agents: MIKSCHE, Gerhard et al.; Aktiebolaget Astra, Patent Department, S-151 85 Södertälje (SE).
(22) International Filing Date: 27 September 1990 (27.09.90)		
(30) Priority data: 8903219-7 2 October 1989 (02.10.89) SE		(81) Designated States: AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH (European patent), CM (OAPI patent), DE (European patent)*, DK (European patent), ES (European patent), FI, FR (European patent), GA (OAPI patent), GB (European patent), HU, IT (European patent), JP, KP, KR, LK, LU (European patent), MC, MG, ML (OAPI patent), MR (OAPI patent), MW, NL (European patent), NO, RO, SD, SE (European patent), SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI patent), US.
(71) Applicant (for all designated States except US): AKTIEBOLAGET ASTRA [SE/SE]; S-151 85 Södertälje (SE).		
(72) Inventors; and		Published
(75) Inventors/Applicants (for US only) : HOFSTRAAT, Robert, Gerrit [NL/NL]; Zwanenveld 25-32, NL-6538 NK Nijmegen (NL). RAIJMAKERS, Petrus, Hendricus [NL/NL]; 11lostraat 19, NL-5402 AA Uden (NL). VRIJHOF, Pieter [NL/NL]; Margrietlaan 10, NL-5351 BX Bergheim (NL).		With international search report.

(54) Title: PROCESS FOR THE MANUFACTURE OF BUDESONIDE



(57) Abstract

The present invention relates to a novel process for the manufacture of (22 R,S)-16α, 17α-butyldenedioxy-11β, 21-dihydroxypregna-1,4-diene-3,20-dione (I) by reacting 11β, 16α, 17α, 21-tetrahydroxypregna-1,4-diene-3,20-dione (II) with butanal, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$ in acetonitrile with p-toluenesulphonic acid as a catalyst.

DESIGNATIONS OF "DE"

Until further notice, any designation of "DE" in any international application whose international filing date is prior to October 3, 1990, shall have effect in the territory of the Federal Republic of Germany with the exception of the territory of the former German Democratic Republic.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MC	Monaco
AU	Australia	FI	Finland	MG	Madagascar
BB	Barbados	FR	France	ML	Mali
BE	Belgium	GA	Gabon	MR	Mauritania
BP	Burkina Fasso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GR	Greece	NL	Netherlands
BJ	Benin	HU	Hungary	NO	Norway
BR	Brazil	IT	Italy	PL	Poland
CA	Canada	JP	Japan	RO	Romania
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
DE	Germany	LJ	Luxembourg	TD	Chad
DK	Denmark			TG	Togo
				US	United States of America

Process for the manufacture of Budesonide


5 Technical field

The present invention relates to a novel process for the manufacture of (22 R,S)-16 α ,17 α -butyridenedioxy-11 β ,21-dihydroxypregna-1,4-diene-3,20-dione (budesonide)

10

15

20




by reacting 11 β ,16 α ,17 α ,21-tetrahydroxypregna-1,4-diene-3,20-dione (16 α -hydroxyprednisolone)

25

30

35



with butanal, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$, in a solvent medium in the presence of an acid catalyst.

Prior art

According to a previously known process disclosed in GB patent no. 1 429 922 Budesonide is manufactured by reacting 16 α -hydroxyprednisolone with butanal in dioxane and with perchloric acid as a catalyst. The product is recovered by diluting the reaction mixture with methylene chloride, and neutralising by washing with aqueous potassium carbonate and water, evaporating the solvent followed by crystallisation from ether/ligroine. The product was further purified by chromatography e.g. on Sephadex.

10 The main disadvantages of dioxane are its skin penetrating and peroxide formation properties. Another disadvantage with this prior art process is perchloric acid, which is a strong oxidizing agent and the use of this catalyst results in a less selective reaction, which in turn makes the subsequent work-up and

15 purification process complicated and expensive.

Disclosure of the invention

The object of the invention is to create a novel process, which gives a more selective reaction and a more simple and economic work-up and purification process.

This is achieved with the process according to the present invention, wherein the reaction is performed in acetonitrile with p-toluenesulphonic acid as a catalyst.

25 The combination of the less basic (compared to dioxane) solvent acetonitrile and the weaker, i.e. non-oxidizing p-toluenesulphonic acid gives a more selective reaction, and also a more simple and economic work-up and purification process compared to the

30 above discussed prior art process using dioxane and perchloric acid.

According to a preferred embodiment of the invention the reaction is stopped by the addition of water and adjustment of the pH of the reaction mixture. This might be done by the addition of sodium hydrogen carbonate in water. The product then crystallizes. The crystals are filtered off, dissolved in methylene chloride and methanol and are then crystallized by the addition

a suitable hydrocarbon, such as ligroine, hexane, cyclohexane or heptane, giving a crude product, which is then recrystallized in methanol/water to give pure budesonide.

- 5 The process according to the invention for the manufacture of budesonide thus consists of two steps.

Step 1. Budesonide crude

10 16α -hydroxyprednisolone is reacted with butanal in acetonitrile. p -Toluenesulphonic acid is added as a catalyst. The reaction mixture is diluted with water and aqueous sodium hydrogen carbonate. After cooling to 5-15°C the crystallized product is filtered off and washed with water. The wet or dried substance is then dissolved in methylene chloride. If the substance used is 15 wet the water phase formed upon dissolution is removed. Methanol is added and the resulting crude budesonide is precipitated by the addition of ligroine or another suitable hydrocarbon (e.g. hexane, heptane or cyclohexane) and is then filtered off.

20 Step 2. Budesonide

The crude budesonide is dissolved in methanol at about 60°C. The solution is filtered through a closed filter and the product is crystallized by the addition of water. After cooling to 5-20°C, filtration and washing with methanol/water the budesonide is 25 dried in vacuum at 40-45°C.

This process is simplified, more economic and less health hazardous compared to prior art processes.

30 Working example

The reaction is carried out in a nitrogen atmosphere. 15.4 g p -toluenesulphonic acid is dissolved into 200 ml acetonitrile. To the solution 50.0 g 16α -hydroxyprednisolone and 17.6 ml butanal are added. The temperature rises to 25°C. After 30 min most of 35 the material is dissolved. Shortly thereafter the product starts to crystallize. After 3 hours the reaction is stopped by the addition of 75 ml aqueous saturated sodium hydrogen carbonate solution, whereupon the product crystallizes. The dried product

is dissolved in methylene chloride and methanol and is crystallized by the addition of ligroine (b.p. 40 - 65), giving crude budesonide.


- 5 The crude budesonide product is recrystallized from methanol/water giving pure budesonide with isomer ratio A:B \approx 1:1 (HPLC), $[\alpha]^{25}_{D}$ 100.0° (c = 0.2; CH_2CL_2); M^+ 430 (theor. 430.5)

claims

1. Process for the manufacture of (22 R,S)-16 α ,17 α -butyldiene-dioxy-11 β ,21-dihydroxypregna-1,4-diene-3,20-dione

5

10




15

by reacting 11 β ,16 α ,17 α ,21-tetrahydroxypregna-1,4-diene-3,20-dione

20

25



30

with butanal, CH₃CH₂CH₂CHO in a solvent medium in the presence of a catalyst, characterised in that the reaction is performed in acetonitrile with p-toluenesulphonic acid as a catalyst.

35

2. Process according to claim 1, characterized in that the reaction is terminated by the addition of water and by adjustment of the pH of the reaction mixture.

3. Process according to claim 1 or 2, characterized in that the crystals obtained upon termination of the reaction are filtered off, dissolved in methylene chloride and methanol and are then crystallized by the addition of a suitable hydrocarbon, such as ligroine, hexane, cyclohexane or heptane, giving a crude product, which is then recrystallized in methanol/water to give pure budesonide.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/SE 90/00619

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all)⁶

According to International Patent Classification (IPC) or to both National Classification and IPC

IPC5: C 07 J 71/00

II. FIELDS SEARCHED

Minimum Documentation Searched⁷

Classification System	Classification Symbols
IPC5	C 07 J

Documentation Searched other than Minimum Documentation
to the Extent that such Documents are Included in Fields Searched⁸

SE,DK,FI,NO classes as above

III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹

Category	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
P,X	US, A, 4925933 (EDIB JAKUPOVIC ET AL.) 15 May 1990, see the whole document --	1-3
X	Chemical Abstracts, volume 106, no. 9, 2 March 1987, (Columbus, Ohio, US), see page 641, abstract 67573q, ES, A, 543211 (Process for the preparation of budesonide) 16 February 1986 --	1-3
X	GB, A, 916996 (OLIN MATHIESON CHEMICAL CORPORATION) 30 January 1963, see especially page 1, lines 45-66 --	1-3
A	EP, A1, 0164636 (SICOR SOCIETA ITALIANA CORTICOSTEROIDI S.P.A.) 18 December 1985, see the whole document -- -----	1-3

* Special categories of cited documents:¹⁰

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step

"Y" document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

IV. CERTIFICATION

Date of the Actual Completion of the International Search

11th December 1990

Date of Mailing of this International Search Report

1990 -12- 21

International Searching Authority

Signature of Authorized Officer

SWEDISH PATENT OFFICE

Anna Sjölund *Anna Sjölund*

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.PCT/SE 90/00619**

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
The members are as contained in the Swedish Patent Office EDP file on **90-11-01**.
The Swedish Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
US-A- 4925933	90-05-15	EP-A-	0262108	88-03-30
		JP-A-	63093795	88-04-25
GB-A- 916996	63-01-30	NONE		
EP-A1- 0164636	85-12-18	JP-A-	61040299	86-02-26
		US-A-	4695625	87-09-22
		US-A-	4835145	89-05-30